

**Basic Facts of GFD +  
Atmospheric LFV, Wind-driven Oceans,  
Paleoclimate & “Tipping Points”**

**Michael Ghil**

**Ecole Normale Supérieure, Paris, and  
University of California, Los Angeles**



**ENS**



*Please visit these sites for more info.*

<https://dept.atmos.ucla.edu/tcd>, <http://www.environnement.ens.fr/>

and [https://www.researchgate.net/profile/Michael\\_Ghil](https://www.researchgate.net/profile/Michael_Ghil)

# Overall Outline

- **Lecture I: Observations and planetary flow theory (GFD<sup>(⌘)</sup>)**
- **Lecture II: Atmospheric LFV<sup>(\*)</sup> & LRF<sup>(\*\*)</sup>**
- **Lecture III: EBMs<sup>(+)</sup>, paleoclimate & “tipping points”**
- ➔ **Lecture IV: Nonlinear & stochastic models—RDS<sup>(⋄)</sup>**
- **Lecture V: Advanced spectral methods—SSA<sup>(±)</sup> *et al.***
- **Lecture VI: The wind-driven ocean circulation**

(⌘) GFD = Geophysical fluid dynamics

(\*) LFV = Low-frequency variability

(\*\*) LRF = Long-range forecasting

(+) EBM = Energy balance model

(⋄) RDS = Random dynamical system

(±) SSA = Singular-spectrum analysis

# Motivation

- The *climate system* is highly *nonlinear and* quite *complex*.
- The system's *major components* — the atmosphere, oceans, ice sheets — *evolve* on many time and space scales.
- Its *predictive understanding* has to rely on the system's physical, chemical and biological modeling, but also on the thorough mathematical analysis of the models thus obtained: *the forest vs. the trees*.
- The *hierarchical modeling* approach allows one to give proper weight to the understanding provided by the models vs. their realism: back-and-forth between *“toy”* (conceptual) and *detailed* (“realistic”) *models*, and between *models* and *data*.
- How do we disentangle *natural variability* from *the anthropogenic forcing*: *can we & should we, or not?*


# Climate and Its Sensitivity

Let's say  $\text{CO}_2$  doubles:

How will “climate” change?

1. Climate is in **stable equilibrium** (fixed point); if so, **mean temperature** will just shift gradually to its new equilibrium value.
2. Climate is **purely periodic**; if so, **mean temperature** will (maybe) shift gradually to its new equilibrium value. But how will the **period, amplitude and phase** of the **limit cycle** change?
3. And how about some “real stuff” now: **chaotic + random**?

Ghil (in *Encycl. Global Environmental Change*, 2002)






# Outline

- The IPCC process: results and uncertainties
- Natural climate variability as a source of uncertainties
  - sensitivity to initial data → error growth
  - sensitivity to model formulation → see below!
- Uncertainties and how to fix them
  - structural stability and other kinds of robustness
  - non-autonomous and random dynamical systems (NDDS & RDS)
- Two illustrative examples
  - the Lorenz convection model
  - an El Niño–Southern Oscillation (ENSO) model
- Linear response theory and climate sensitivity
- Conclusions and references
  - natural variability and anthropogenic forcing: the “grand unification”
  - selected bibliography


# CO2 IN THE ATMOSPHERE



# Temperatures and GHGs

Greenhouse gases (GHGs) go up,  
temperatures go up:

It's gotta do with us, at least a bit,  
doesn't it?



Wikicommons, from  
Hansen *et al.* (*PNAS*, 2006);  
see also <http://data.giss.nasa.gov/gistemp/graphs/>

# Unfortunately, things aren't all that easy!

**What to do?**

**Try to achieve better interpretation of, and agreement between, models ...**

Ghil, M., 2002: Natural climate variability, in *Encyclopedia of Global Environmental Change*, T. Munn (Ed.), Vol. 1, Wiley

Natural variability introduces additional complexity into the anthropogenic climate change problem


The most common interpretation of observations and GCM simulations of climate change is still in terms of a scalar, linear Ordinary Differential Equation (ODE)

$$c \frac{dT}{dt} = -kT + Q$$

$k = \sum k_i$  – feedbacks (+ve and -ve)

$Q = \sum Q_j$  – sources & sinks

$Q_j = Q_j(t)$



Linear response to CO<sub>2</sub> vs. observed change in T

Hence, we need to consider instead a system of nonlinear Partial Differential Equations (PDEs), with parameters and multiplicative, as well as additive forcing (deterministic + stochastic)

$$\frac{dX}{dt} = N(X, t, \mu, \beta)$$

# Global warming and its socio-economic impacts

Temperatures rise:

- What about impacts?
- How to adapt?

The answer, my friend, is blowing in the wind, *i.e.*, it depends on the accuracy and reliability of the forecast ...

Source : IPCC (2007),  
AR4, WGI, SPM




Figure SPM.5. Solid lines are multi-model global averages of surface warming (relative to 1980–1999) for the scenarios A2, A1B and B1, shown as continuations of the 20th century simulations. Shading denotes the  $\pm 1$  standard deviation range of individual model annual averages. The orange line is for the experiment where concentrations were held constant at year 2000 values. The grey bars at right indicate the best estimate (solid line within each bar) and the likely range assessed for the six SRES marker scenarios. The assessment of the best estimate and likely ranges in the grey bars includes the AOGCMs in the left part of the figure, as well as results from a hierarchy of independent models and observational constraints. (Figures 10.4 and 10.29)

# Global warming and its socio-economic impacts– II


Temperatures rise:

- What about impacts?
- How to adapt?

## AR5 vs. AR4

A certain air of *déjà vu*:  
GHG “scenarios” have been replaced by “representative concentration pathways” (RCPs), more dire predictions, but the **uncertainties** remain.


Source : IPCC (2013),  
AR5, WGI, SPM





# Outline

- The IPCC process: results and uncertainties
- **Natural climate variability as a source of uncertainties**
  - sensitivity to initial data → error growth
  - sensitivity to model formulation → see below!
- Uncertainties and how to fix them
  - structural stability and other kinds of robustness
  - non-autonomous and random dynamical systems (NDDS & RDS)
- Two illustrative examples
  - the Lorenz convection model
  - an El Niño–Southern Oscillation (ENSO) model
- Linear response theory and climate sensitivity
- Conclusions and references
  - natural variability and anthropogenic forcing: the “grand unification”
  - selected bibliography



Courtesy Tim Palmer, 2009




# Exponential divergence vs. “coarse graining”

The classical view of dynamical systems theory is:  
positive Lyapunov exponent →  
trajectories diverge exponentially

But the presence of multiple regimes implies a much more structured behavior in phase space

Still, the probability distribution function (pdf), when calculated forward in time, is pretty smeared out

L. A. Smith (*Encycl. Atmos. Sci.*, 2003)



# So what's it gonna be like, by 2100?

**Table SPM.2.** Recent trends, assessment of human influence on the trend and projections for extreme weather events for which there is an observed late-20th century trend. (Tables 3.7, 3.8, 9.4; Sections 3.8, 5.5, 9.7, 11.2–11.9)


| Phenomenon <sup>a</sup> and direction of trend   | Likelihood that trend occurred in late 20th century (typically post 1960) | Likelihood of a human contribution to observed trend <sup>b</sup> | Likelihood of future trends based on projections for 21st century using SRES scenarios |
|--|---|---|--|
| Warmer and fewer cold days and nights over most land areas   | <i>Very likely<sup>c</sup></i>  | <i>Likely<sup>d</sup></i>   | <i>Virtually certain<sup>d</sup></i>   |
| Warmer and more frequent hot days and nights over most land areas  | <i>Very likely<sup>e</sup></i>  | <i>Likely (nights)<sup>d</sup></i>                                | <i>Virtually certain<sup>d</sup></i>   |
| Warm spells/heat waves. Frequency increases over most land areas   | <i>Likely</i>   | <i>More likely than not<sup>f</sup></i>                           | <i>Very likely</i>   |
| Heavy precipitation events. Frequency (or proportion of total rainfall from heavy falls) increases over most areas | <i>Likely</i>   | <i>More likely than not<sup>f</sup></i>                           | <i>Very likely</i>   |
| Area affected by droughts increases  | <i>Likely in many regions since 1970s</i>                                 | <i>More likely than not</i>                                       | <i>Likely</i>  |
| Intense tropical cyclone activity increases  | <i>Likely in some regions since 1970</i>                                  | <i>More likely than not<sup>f</sup></i>                           | <i>Likely</i>  |
| Increased incidence of extreme high sea level (excludes tsunamis) <sup>g</sup>                                     | <i>Likely</i>   | <i>More likely than not<sup>h</sup></i>                           | <i>Likely<sup>i</sup></i>  |

# Outline

- The IPCC process: results and uncertainties
- Natural climate variability as a source of uncertainties
  - sensitivity to initial data → error growth
  - sensitivity to model formulation → see below!
- **Uncertainties and how to fix them**
  - structural stability and other kinds of robustness
  - non-autonomous and random dynamical systems (NDDS & RDS)
- Two illustrative examples
  - the Lorenz convection model
  - an El Niño–Southern Oscillation (ENSO) model
- Linear response theory and climate sensitivity
- Conclusions and references
  - natural variability and anthropogenic forcing: the “grand unification”
  - selected bibliography

# How important are different sources of uncertainty?

- Varies, but typically no single source dominates.



Uncertainties in winter precipitation changes for the 2080s relative to 1961-90, at a 25km box in SE England

Source: Met Office



# Can we, nonlinear dynamicists, help?

The uncertainties  
might be *intrinsic*,  
rather than mere  
“tuning problems”

If so, maybe  
*stochastic structural  
stability* could help!

Might fit in nicely with  
recent taste for  
“stochastic  
parameterizations”




Figure 7.5-1. The three towers of differentiable dynamics.

*The DDS dream of structural stability* (from Abraham & Marsden, 1978)

# Non-autonomous Dynamical Systems

## A linear, dissipative, forced example: *forward vs. pullback attraction*

Consider the scalar, linear ordinary differential equation (ODE)

$$\dot{x} = -\alpha x + \sigma t, \quad \alpha > 0, \quad \sigma > 0.$$


The autonomous part of this ODE,  $\dot{x} = -\alpha x$ , is **dissipative** and all solutions  $x(t; x_0) = x(t; x(0) = x_0)$  converge to 0 as  $t \rightarrow +\infty$ .

What about the non-autonomous, forced ODE? As the energy being put into the system by the forcing is dissipated, we expect things to change in time. In fact, if we “pull back” far enough, replace  $x(t; x_0)$  by  $x(s, t; x_0) = x(s, t; x(s) = x_0)$ ,

$x(s, t; x_0)$ , with  $x_0$  varying

and let  $s \rightarrow -\infty$ , we get the **pullback attractor**  $a = a(t)$  in the figure,

$$a(t) = \frac{\sigma}{\alpha} \left( t - \frac{1}{\alpha} \right).$$



## Remarks

- We've just shown that:

$$|x(t, s; x_0) - a(t)| \xrightarrow{s \rightarrow -\infty} 0 ; \text{ for every } t \text{ fixed,}$$

and for all initial data  $x_0$ , with  $a(t) = \frac{\sigma}{\alpha}(t - 1/\alpha)$ .

- We've just encountered the concept of **pullback attraction**; here  $\{a(t)\}$  is the **pullback attractor** of the system (1).
- What does it mean physically?

The pullback attractor provides a way to assess an **asymptotic regime at time  $t$**  — the time at which we observe the system — for a system starting to evolve from the remote past  $s$ ,  $s \ll t$ .

- This asymptotic regime evolves with time: it is a dynamical object.
- Dissipation now leads to a dynamic object rather than to a static one, like the strange attractor of an autonomous system.

## Remarks

- We've just shown that:

$$|x(t, s; x_0) - a(t)| \xrightarrow{s \rightarrow -\infty} 0 ; \text{ for every } t \text{ fixed,}$$

and for all initial data  $x_0$ , with  $a(t) = \frac{\sigma}{\alpha}(t - 1/\alpha)$ .

- We've just encountered the concept of **pullback attraction**; here  $\{a(t)\}$  is the **pullback attractor** of the system (1).
- What does it mean physically?  
The pullback attractor provides a way to assess an **asymptotic regime at time  $t$**  — the time at which we observe the system — for a system starting to evolve from the remote past  $s$ ,  $s \ll t$ .
- This asymptotic regime evolves with time: it is a dynamical object.
- Dissipation now leads to a dynamic object rather than to a static one, like the strange attractor of an autonomous system.




# A little history of climate & stochasticity

- A. Einstein's (1905) Brownian motion paper.
- K. Itō (prof. at Kyoto U., RIMS director) formulates Itō calculus in 1942, enables solution of stochastic differential equations (SDEs); Itō's lemma is the stochastic counterpart of Leibniz's chain rule for differentiation.
- K. Hasselmann (*Tellus*, 1976) describes climate as Brownian motion, with weather the stochastic driver.
- In this view, the deterministic part of the model is stable, and random perturbations decay to the mean.



Kiyoshi Itō



Auto-regressive (AR) decay

# Random Dynamical Systems (RDS), I - RDS theory

This theory is the counterpart for randomly forced dynamical systems (RDS) of the *geometric theory* of ordinary differential equations (ODEs). It allows one to treat stochastic differential equations (SDEs) — and more general systems driven by noise — as flows in (phase space)  $\times$  (probability space).

SDE  $\sim$  ODE, RDS  $\sim$  DDS, L. Arnold (1998)  $\sim$  V.I. Arnol'd (1983).

## Setting:

- (i) A phase space  $X$ . **Example:**  $\mathbb{R}^n$ .
- (ii) A probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ . **Example:** The Wiener space  $\Omega = C_0(\mathbb{R}; \mathbb{R}^n)$  with Wiener measure  $\mathbb{P}$ .
- (iii) A model of the noise  $\theta(t) : \Omega \rightarrow \Omega$  that preserves the measure  $\mathbb{P}$ , i.e.  $\theta(t)\mathbb{P} = \mathbb{P}$ ;  $\theta$  is called **the driving system**.  
**Example:**  $W(t, \theta(s)\omega) = W(t + s, \omega) - W(s, \omega)$ ;  
it starts the noise at  $s$  instead of  $t = 0$ .
- (iv) A mapping  $\varphi : \mathbb{R} \times \Omega \times X \rightarrow X$  with the cocycle property.  
**Example:** The solution operator of an SDE.

# Random Dynamical Systems (RDS), I - RDS theory

This theory is the counterpart for randomly forced dynamical systems (RDS) of the *geometric theory* of ordinary differential equations (ODEs). It allows one to treat stochastic differential equations (SDEs) — and more general systems driven by noise — as flows in (phase space)  $\times$  (probability space).

SDE  $\sim$  ODE, RDS  $\sim$  DDS, L. Arnold (1998)  $\sim$  V.I. Arnol'd (1983).

## Setting:


- (i) A phase space  $X$ . **Example:**  $\mathbb{R}^n$ .
- (ii) A probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ . **Example:** The Wiener space  $\Omega = C_0(\mathbb{R}; \mathbb{R}^n)$  with Wiener measure  $\mathbb{P}$ .
- (iii) A model of the noise  $\theta(t) : \Omega \rightarrow \Omega$  that preserves the measure  $\mathbb{P}$ , i.e.  $\theta(t)\mathbb{P} = \mathbb{P}$ ;  $\theta$  is called **the driving system**.  
**Example:**  $W(t, \theta(s)\omega) = W(t + s, \omega) - W(s, \omega)$ ;  
it starts the noise at  $s$  instead of  $t = 0$ .
- (iv) A mapping  $\varphi : \mathbb{R} \times \Omega \times X \rightarrow X$  with the cocycle property.  
**Example:** The solution operator of an SDE.

# RDS, III- Random attractors (RAs)

A random attractor  $\mathcal{A}(\omega)$  is both *invariant* and “pullback” *attracting*:

- (a) **Invariant:**  $\varphi(t, \omega)\mathcal{A}(\omega) = \mathcal{A}(\theta(t)\omega)$ .
- (b) **Attracting:**  $\forall B \subset X, \lim_{t \rightarrow \infty} \text{dist}(\varphi(t, \theta(-t)\omega)B, \mathcal{A}(\omega)) = 0$  a.s.

*Pullback attraction to  $\mathcal{A}(\omega)$*




# Outline

- The IPCC process: results and uncertainties
- Natural climate variability as a source of uncertainties
  - sensitivity to initial data → error growth
  - sensitivity to model formulation → see below!
- Uncertainties and how to fix them
  - structural stability and other kinds of robustness
  - non-autonomous and random dynamical systems (NDDS & RDS)
- **Two illustrative examples**
  - the Lorenz convection model
  - an El Niño–Southern Oscillation (ENSO) model
- Linear response theory and climate sensitivity
- Conclusions and references
  - natural variability and anthropogenic forcing: the “grand unification”
  - selected bibliography


# Random attractor of the stochastic Lorenz system

## Snapshot of the random attractor (RA)




- A **snapshot** of the RA,  $\mathcal{A}(\omega)$ , computed at a fixed time  $t$  and for the **same realization**  $\omega$ ; it is made up of points transported by the stochastic flow, from the remote past  $t - T$ ,  $T \gg 1$ .
- We use **multiplicative noise** in the deterministic Lorenz model, with the classical parameter values  $b = 8/3$ ,  $\sigma = 10$ , and  $r = 28$ .
- Even computed **pathwise**, this object supports meaningful **statistics**.

# Sample measures supported by the R.A.



- We compute the probability measure on the R.A. at some fixed time  $t$ , and for a fixed realization  $\omega$ . We show a “projection”,  $\int \mu_\omega(x, y, z) dy$ , with **multiplicative noise**:  $dx_i = \text{Lorenz}(x_1, x_2, x_3) dt + \alpha x_i dW_t; i \in \{1, 2, 3\}$ .
- **10 million of initial points** have been used for this picture!

# Sample measure supported by the R.A.




- Still 1 Billion I.D., and  $\alpha = 0.5$ . Another one?



## Sample measures evolve with time.

- Recall that these sample measures are the **frozen statistics** at a time  $t$  for a realization  $\omega$ .
- How do these **frozen statistics** evolve with time?
- **Action!**



A day in the life of the Lorenz (1963) model's random attractor, or LORA for short;  
see SI in Chekroun, Simonnet & Ghil (2011, *Physica D*) or  
Vimeo movie: <https://vimeo.com/240039610>

# Outline

- The IPCC process: results and uncertainties
- Natural climate variability as a source of uncertainties
  - sensitivity to initial data → error growth
  - sensitivity to model formulation → see below!
- Uncertainties and how to fix them
  - structural stability and other kinds of robustness
  - non-autonomous and random dynamical systems (NDDS & RDS)
- Two illustrative examples
  - the Lorenz convection model
  - an El Niño–Southern Oscillation (ENSO) model
- **Linear response theory and climate sensitivity**
- Conclusions and references
  - natural variability and anthropogenic forcing: the “grand unification”
  - selected bibliography


# Climate and Its Sensitivity

Let's say  $\text{CO}_2$  doubles:

How will “climate” change?

1. Climate is in **stable equilibrium** (fixed point); if so, **mean temperature** will just shift gradually to its new equilibrium value.
2. Climate is **purely periodic**; if so, **mean temperature** will (maybe) shift gradually to its new equilibrium value. But how will the **period, amplitude and phase** of the **limit cycle** change?
3. And how about some “real stuff” now: **chaotic + random**?

Ghil (in *Encycl. Global Environmental Change*, 2002)




# Classical Strange Attractor

Physically **closed** system, modeled mathematically as **autonomous** system: neither deterministic (anthropogenic) nor random (natural) forcing.

The **attractor** is **strange**, but still fixed in time ~ “**irrational**” number.

**Climate sensitivity** ~ change in the **average value (first moment)** of the coordinates  $(x, y, z)$  as a **parameter  $\lambda$**  changes.




# Random Attractor

Physically **open** system, modeled mathematically as **non-autonomous** system: allows for deterministic (anthropogenic) as well as random (natural) forcing.

The **attractor** is “**pullback**” and evolves in time  $\sim$  “**imaginary**” or “**complex**” number.


**Climate sensitivity**  $\sim$  change in the statistical properties (first and **higher-order moments**) of the **attractor** as one or more parameters ( $\lambda$ ,  $\mu$ , ...) change.

Ghil (*Encyclopedia of Atmospheric Sciences*, 2<sup>nd</sup> ed., 2012)





# Spatio-temporal evolution of ENSO episode




# Parameter dependence – I

$$\delta = 0.9557$$


It can be smooth or it can be rough:  
Niño-3 SSTs from intermediate coupled model  
for ENSO (Jin, Neelin & Ghil, 1994, 1996)

Skewness & kurtosis of the SSTs:  
time series of 4000 years,


$$\Delta\delta = 3 \cdot 10^{-4}$$



Skewness dependence



Kurtosis dependence



M. Chekroun (work in progress)



# Sample measures for an NDDE model of ENSO

## The Galanti-Tziperman (GT) model (JAS, 1999)


$$\frac{dT}{dt} = -\epsilon_T T(t) - M_0(T(t) - T_{sub}(h(t))),$$

Neutral delay-differential equation (NDDE), derived from Cane-Zebiak and Jin-Neelin models for ENSO:  $T$  is East-basin SST and  $h$  is thermocline depth.

$$h(t) = M_1 e^{-\epsilon_m(\tau_1 + \tau_2)} h(t - \tau_1 - \tau_2) - M_2 \tau_1 e^{-\epsilon_m(\frac{\tau_1}{2} + \tau_2)} \mu(t - \tau_2 - \frac{\tau_1}{2}) T(t - \tau_2 - \frac{\tau_1}{2}) + M_3 \tau_2 e^{-\epsilon_m \frac{\tau_2}{2}} \mu(t - \frac{\tau_2}{2}) T(t - \frac{\tau_2}{2}).$$

Seasonal forcing given by  $\mu(t) = 1 + \epsilon \cos(\omega t + \phi)$ .  
 The pullback attractor and its invariant measures were computed.

Figure shows the changes in the mean, 2<sup>nd</sup> & 4<sup>th</sup> moment of  $h(t)$ , along with the Wasserstein distance  $d_W$ , for changes of 0–5% in the delay parameter  $\tau_{K,0}$ .



Note intervals of both **smooth** & **rough** dependence!

# How to define climate sensitivity or, What happens when there's natural variability?


One usually defines **climate sensitivity**  $\gamma$  as  $\Delta T/\Delta Q$ ,  
where  $T$  is global mean temperature in  $^{\circ}\text{C}$ ,  
and  $\Delta Q$  is insolation change in %.  
Thus  $\gamma \approx 1^{\circ}\text{C}$  per 1 % change in  $Q$ .

But there is much more to climate than mean  $T$ : there is the actual distribution of temperatures in time and space, there's extrema of temperatures and of precipitations, etc., as in the figure.

So we would like a better, more flexible definition, which does take into account these “details,” as well as chaotic behavior:



$$\gamma = \partial d_W / \partial \tau \text{ or}$$

$$\gamma = \Delta d_W / \Delta \tau$$




# How to define climate sensitivity or, What happens when there's natural variability?

This definition allows us to watch how “the earth moves,” as it is affected by both natural and anthropogenic forcing, in the presence of natural variability, which includes both chaotic & random behavior:





$$\gamma = \partial d_W / \partial \tau \text{ or}$$
$$\gamma = \Delta d_W / \Delta \tau$$



# Parameter dependence – II


When it is smooth, one can optimize a GCM's single-parameter dependence



ICTP AGCM (Neelin, Bracco, Luo, McWilliams & Meyerson, *PNAS*, 2010)

# Parameter dependence – III

Multi-objective algorithms avoid arbitrary weighting of criteria in a unique cost function:



Optimization algorithms that are  $\mathcal{O}(N)$  and  $\mathcal{O}(N^2)$ , rather than  $\mathcal{O}(S^N)$ , where  $N$  is the number of parameters and  $S$  is the sampling density.

ICTP AGCM (Neelin, Bracco, Luo, McWilliams & Meyerson, *PNAS*, 2010)

# Outline

- The IPCC process: results and uncertainties
- Natural climate variability as a source of uncertainties
  - sensitivity to initial data → error growth
  - sensitivity to model formulation → see below!
- Uncertainties and how to fix them
  - structural stability and other kinds of robustness
  - non-autonomous and random dynamical systems (NDDS & RDS)
- Two illustrative examples
  - the Lorenz convection model
  - an El Niño–Southern Oscillation (ENSO) model
- Linear response theory and climate sensitivity
- **Conclusions and references**
  - natural variability and anthropogenic forcing: the “grand unification”
  - selected bibliography

# Concluding remarks, I – RDS and RAs

## *Summary*

- A change of paradigm from **closed, autonomous systems** to **open, non-autonomous ones**.
- Random attractors are (i) spectacular, (ii) useful, and (iii) just starting to be explored for climate applications.

## *Work in progress*

- Study the effect of specific **stochastic parametrizations** on model robustness.
- Applications to **intermediate models and GCMs**.
- Implications for **climate sensitivity**.
- Implications for **predictability?**

# Yet another (grand?) unification

Lorenz (*JAS*, 1963)

Climate is deterministic and autonomous,  
but highly nonlinear.

Trajectories diverge exponentially,  
forward asymptotic PDF is multimodal.

Hasselmann (*Tellus*, 1976)


Climate is stochastic and noise-driven,  
but quite linear.

Trajectories decay back to the mean,  
forward asymptotic PDF is unimodal.


**Grand unification (?)**

Climate is deterministic + stochastic,  
as well as highly nonlinear.


Internal variability and forcing interact  
strongly, **change and sensitivity**  
refer to both mean and higher moments.



$X(t, \omega; X_0)$ , with  $X_0$  varying



Time-dependent invariant measure of the GT-model





# Concluding remarks, II – Climate change & climate sensitivity

## *What do we know?*

- It's getting warmer.
- We do contribute to it.
- So we should act as best we know and can!

## *What do we know less well?*

- By how much?
  - Is it getting warmer ...
  - Do we contribute to it ...
- How does the climate system (atmosphere, ocean, ice, etc.) really work?
- How does natural variability interact with anthropogenic forcing?

## *What to do?*

- Better understand the system and its forcings.
- Explore the models', and the system's, robustness and sensitivity
  - stochastic structural and statistical stability!
  - linear response = response function + susceptibility function!!

# Concluding remarks, II – Climate change & climate sensitivity

## *What do we know?*

- It's getting warmer.
- We do contribute to it.
- So we should act as best we know and can!

## *What do we know less well?*

- By how much?
  - Is it getting warmer ...
  - Do we contribute to it ...
- How does the climate system (atmosphere, ocean, ice, etc.) really work?
- How does natural variability interact with anthropogenic forcing?

## *What to do?*

- Better understand the system and its forcings.
- Explore the models', and the system's, robustness and sensitivity
  - stochastic structural and statistical stability!
  - linear response = response function + susceptibility function!!

# Concluding remarks, II – Climate change & climate sensitivity

## *What do we know?*

- It's getting warmer.
- We do contribute to it.
- So we should act as best we know and can!

## *What do we know less well?*

- By how much?
  - Is it getting warmer ...
  - Do we contribute to it ...
- How does the climate system (atmosphere, ocean, ice, etc.) really work?
- How does natural variability interact with anthropogenic forcing?

## *What to do?*

- Better understand the system and its forcings.
- Explore the models', and the system's, **robustness and sensitivity**
  - **stochastic structural and statistical stability!**
  - **linear response = response function + susceptibility function!!**

# Some general references

- Andronov, A.A., and L.S. Pontryagin, 1937: Systèmes grossiers. *Dokl. Akad. Nauk. SSSR*, **14**(5), 247–250.
- Arnold, L., 1998: *Random Dynamical Systems*, Springer Monographs in Math., Springer, 625 pp.
- Arnol'd, V. I., 1983: *Geometrical Methods in the Theory of Ordinary Differential Equations*, Springer-Verlag, New York/Heidelberg/Berlin, 334 pp.
- Charney, J.G., *et al.*, 1979: *Carbon Dioxide and Climate: A Scientific Assessment*. National Academic Press, Washington, D.C.
- Chekroun, M. D., E. Simonnet, and M. Ghil, 2011: Stochastic climate dynamics: Random attractors and time-dependent invariant measures, *Physica D*, **240**, 1685–1700.
- Ghil, M., R. Benzi, and G. Parisi (Eds.), 1985: *Turbulence and Predictability in Geophysical Fluid Dynamics and Climate Dynamics*, North-Holland, 449 pp.
- Ghil, M., and S. Childress, 1987: *Topics in Geophysical Fluid Dynamics: Atmospheric Dynamics, Dynamo Theory and Climate Dynamics*, Ch. 5, Springer-Verlag, New York, 485 pp.
- Ghil, M., M.D. Chekroun, and E. Simonnet, 2008: Climate dynamics and fluid mechanics: Natural variability and related uncertainties, *Physica D*, **237**, 2111–2126.
- Houghton, J.T., G.J. Jenkins, and J.J. Ephraums (Eds.), 1991: *Climate Change, The IPCC Scientific Assessment*, Cambridge Univ. Press, Cambridge, MA, 365 pp.
- Lorenz, E.N., 1963: Deterministic nonperiodic flow. *J. Atmos. Sci.*, **20**, 130–141.
- Neelin, J.D., A. Bracco *et al.*, 2010: Considerations for parameter optimization and sensitivity in climate models, *Proc. Natl. Acad. Sci. USA*, **107**, 21,349–21,354.
- Ruelle, D., 1997: Application of hyperbolic dynamics to physics: Some problems and conjectures, *Bull. Amer. Math. Soc.*, **41**, 275–278.
- Solomon, S., *et al.* (Eds.). *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the IPCC*, Cambridge Univ. Press, 2007.

# The Lorenz (1963a) convection model

**Problem 2:** Find the appropriate software to compute the Lorenz “butterfly” and use it to do so.

**Problem 4:** Find the appropriate software to compute the statistics of the Lorenz “butterfly” – e.g., pdf, EOFs – and use it to do so.

## *Glossary*

pdf = probability density function

EOF = empirical orthogonal function

**Problem 8:** Add some noise to the Lorenz convection model and compute:

- a) some sample solutions;
- b) the invariant measure (more precisely, an approximate pdf);
- c) the random attractor; and
- d) its sensitivity to parameter changes.

**Reserve slides**



# Galileo on math, science & opinions, 1

La filosofia è scritta in questo grandissimo libro che continuamente ci sta aperto innanzi a gli occhi (io dico l'universo), ma non si può intendere se prima non s'impara a intender la lingua, e conoscer i caratteri, ne' quali è scritto. Egli è scritto in lingua matematica, e i caratteri son triangoli, cerchi ed altre figure geometriche, senza i quali mezzi è impossibile a intenderne umanamente parola; senza questi è un aggirarsi vanamente per un oscuro laberinto.

G. Galilei, *Il Saggiatore*, VI, 232)

Philosophy is written in this grand book — I mean the Universe—which stands continually open to our gaze, but it cannot be understood unless one first learns to comprehend the language and interpret the characters in which it is written. It is written in the language of mathematics, and its characters are triangles, circles and other geometrical figures, without which it is humanly impossible to understand a single word of it.

## letters to nature

*Nature* 350, 324 - 327 (1991); doi:10.1038/350324a0

# Interdecadal oscillations and the warming trend in global temperature time series

M. Ghil & R. Vautard

THE ability to distinguish a warming trend from natural variability is critical for an understanding of the climatic response to increasing greenhouse-gas concentrations. Here we use singular spectrum analysis<sup>1</sup> to analyse the time series of global surface air temperatures for the past 135 years<sup>2</sup>, allowing a secular warming trend and a small number of oscillatory modes to be separated from the noise. The trend is flat until 1910, with an increase of 0.4 °C since then. The oscillations exhibit interdecadal periods of 21 and 16 years, and interannual periods of 6 and 5 years. The interannual oscillations are probably related to global aspects of the El Niño-Southern Oscillation (ENSO) phenomenon<sup>3</sup>. The interdecadal oscillations could be associated with changes in the extratropical ocean circulation<sup>4</sup>. The oscillatory components have combined (peak-to-peak) amplitudes of 0.2 °C, and therefore limit our ability to predict whether the inferred secular warming trend of 0.005 °Cyr<sup>-1</sup> will continue. This could postpone incontrovertible detection of the greenhouse warming signal for one or two decades.



## Galileo on math, science & opinions, 2

*Sì perché l'autorità dell'opinione di mille nelle scienze non val per una scintilla di ragione di un solo.*

In questions of science, the authority of a thousand is not worth the humble reasoning of a single individual.

Galileo Galilei, *Venere, Luna e Pianeti Medicei,*  
*e nuove apparenze di Saturno*, p. 8/20

# Applications to a nonlinear stochastic El Niño model

Chekroun, Simonnet and Ghil, 2011

Timmerman & Jin (*Geophys. Res. Lett.*, 2002) have derived the following low-order, tropical-atmosphere–ocean model. The model has three variables: thermocline depth anomaly  $h$ , and SSTs  $T_1$  and  $T_2$  in the western and eastern basin.

$$\begin{aligned}\dot{T}_1 &= -\alpha(T_1 - T_r) - \frac{2\varepsilon u}{L}(T_2 - T_1), \\ \dot{T}_2 &= -\alpha(T_2 - T_r) - \frac{w}{H_m}(T_2 - T_{sub}), \\ \dot{h} &= r(-h - bL\tau/2).\end{aligned}$$

The related diagnostic equations are:

$$\begin{aligned}T_{sub} &= T_r - \frac{T_r - T_{r0}}{2} [1 - \tanh(H + h_2 - z_0)/h^*] \\ \tau &= \frac{a}{\beta} (T_1 - T_2) [\xi_t - 1].\end{aligned}$$


- $\tau$ : the wind stress anomalies,  $w = -\beta\tau/H_m$ : the equatorial upwelling.
- $u = \beta L\tau/2$ : the zonal advection,  $T_{sub}$ : the subsurface temperature.

**Wind stress bursts** are modeled as white noise  $\xi_t$  of variance  $\sigma$ , and  $\varepsilon$  measures the strength of the **zonal advection**.




# Random attractors: the frozen statistics

## Random Shil'nikov horseshoes




$$\sigma=0.005$$



$$\sigma=0.05$$


- Horseshoes can be noise-excited, left: a weakly-perturbed limit cycle, right: the same with larger noise.
- Golden: most frequently-visited areas; white 'plus' sign: most visited.

# An episode in the random's attractor life






# Atmospheric CO<sub>2</sub> at Mauna Loa Observatory



# GHGs rise!

It's gotta do with us, at least a bit, ain't it?  
But just how much?

RADIATIVE FORCING COMPONENTS



©IPCC 2007: WG1-AR4


IPCC (2007)



*AR4 adjustment of 20<sup>th</sup> century simulation*

[www.lseca](http://www.lseca)


Hindcasts and Forecasts of Global Mean Temperature



Ed Tredger  
(PhD thesis, LSE, 2009)



# RDS, II - A Geometric View of SDEs



- $\varphi$  is a random dynamical system (RDS)
- $\Theta(t)(x, \omega) = (\theta(t)\omega, \varphi(t, \omega)x)$  is a flow on the bundle

# Non-autonomous Dynamical Systems - I

## A linear example as a paradigm

Let us first start with a **very difficult problem**:

$$\text{Study the "dynamics" of } \dot{x} = -\alpha x + \sigma t, \quad \alpha, \sigma > 0. \quad (1)$$

### First remarks:

- The system  $\dot{x} = -\alpha x$ , i.e. the autonomous part of (1), is **dissipative**. All the solutions of  $\dot{x} = -\alpha x$  converge to 0 as  $t \rightarrow +\infty$ .
- Is it the case for (1)? Certainly not!  
The **autonomous part** is **forced**; we even introduce an **infinite energy** over an infinite time interval:  $\int_0^{+\infty} t \, dt = +\infty!$   
Forward attraction seems to be ill adapted to time-dependent forcing.

### Goal:

Find a concept of attraction that is:

- (i) compatible with the forward concept, when there is no forcing; and
- (ii) provides a way to assess the effect of dissipation in some sense.

For that let's do some computations...

# Non-autonomous Dynamical Systems - I

## A linear example as a paradigm

Let us first start with a **very difficult problem**:

$$\text{Study the "dynamics" of } \dot{x} = -\alpha x + \sigma t, \quad \alpha, \sigma > 0. \quad (1)$$

### First remarks:

- The system  $\dot{x} = -\alpha x$ , i.e. the autonomous part of (1), is **dissipative**. All the solutions of  $\dot{x} = -\alpha x$  converge to 0 as  $t \rightarrow +\infty$ .
- Is it the case for (1)? Certainly not!  
The **autonomous part** is **forced**; we even introduce an **infinite energy** over an infinite time interval:  $\int_0^{+\infty} t \, dt = +\infty!$   
Forward attraction seems to be ill adapted to time-dependent forcing.

### Goal:

Find a concept of attraction that is:

- (i) compatible with the forward concept, when there is no forcing; and
- (ii) provides a way to assess the effect of dissipation in some sense.

For that let's do some computations...



# Non-autonomous Dynamical Systems - I

## A linear example as a paradigm

Let us first start with a **very difficult problem**:

$$\text{Study the "dynamics" of } \dot{x} = -\alpha x + \sigma t, \quad \alpha, \sigma > 0. \quad (1)$$

### First remarks:

- The system  $\dot{x} = -\alpha x$ , i.e. the autonomous part of (1), is **dissipative**. All the solutions of  $\dot{x} = -\alpha x$  converge to 0 as  $t \rightarrow +\infty$ .
- Is it the case for (1)? Certainly not!  
The **autonomous part** is **forced**; we even introduce an **infinite energy** over an infinite time interval:  $\int_0^{+\infty} t \, dt = +\infty!$   
Forward attraction seems to be ill adapted to time-dependent forcing.

### Goal:

Find a concept of attraction that is:


- (i) compatible with the forward concept, when there is no forcing; and
- (ii) provides a way to assess the effect of dissipation in some sense.

**For that let's do some computations...**


# A French garden near the castle of La Roche-Guyon



# Devil's quarry for a coupling parameter $\varepsilon = 0.15$ : a web of resonances




# Effect of the noise on Devil's quarry




# Devil's Bleachers in a 1-D ENSO Model

Ratio of ENSO frequency to annual cycle




# Sample measure supported by the R.A.



- 1 Billion I.D., and a different color palette!
- Intensity is  $\alpha = 0.2$ .
- Do you want different noise intensities?

# Sample measure supported by the R.A.


Another proj. of the sample measure, "friendlier"



- The next slides are similar, with different **noise level  $\alpha$**  and more I.D....



# Sample measure supported by the R.A.



- Here  $\alpha = 0.4$ . The sample measure is approximated for another realization  $\omega$  of the noise, starting from 8 billion I.D.
- Now more serious stuff is coming...

## Sample measures evolve with time.

- Recall that these sample measures are the **frozen statistics** at a time  $t$  for a realization  $\omega$ .
- How do these **frozen statistics** evolve with time?
- **Action!**

## Sample measures evolve with time.

- Recall that these sample measures are the **frozen statistics** at a time  $t$  for a realization  $\omega$ .
- How do these **frozen statistics** evolve with time?
- Action!

## The Sinai-Ruelle-Bowen (SRB) property

- RDS theory offers a rigorous way to define random versions of stable and unstable manifolds, via the Lyapunov spectrum, the Oseledec multiplicative theorem, and a random version of the Hartman-Grobman theorem.
- When the sample measures  $\mu_\omega$  of an RDS have **absolutely continuous conditional measures** on the random unstable manifolds, then  $\mu_\omega$  is called a **random SRB measure**.
- If the sample measure of an RDS  $\varphi$  is SRB, then its a “physical” measure in the sense that:

$$\lim_{s \rightarrow -\infty} \frac{1}{t-s} \int_s^t G \circ \varphi(s, \theta_{-s}\omega)x \, ds = \int_{\mathcal{A}(\theta_t\omega)} G(x) \mu_{\theta_t\omega}(dx), \quad (3)$$

for almost every  $x \in X$  (in the Lebesgue sense), and for every continuous **observable**  $G : X \rightarrow \mathbb{R}$ .

- The measure  $\mu_\omega$  is also the image of the Lebesgue measure under the stochastic flow  $\varphi$ : for each region of  $\mathcal{A}(\omega)$ , it gives the **probability to end up** on that region, when starting from a volume.

## A remarkable theorem of Ledrappier and Young (1988)

- Ledrappier and Young have proved that, that if the **stationary solution**,  $\rho$ , of the **Fokker-Planck equation** associated to an SDE presenting a Lyapunov exponent  $> 0$ , has a **density** w.r.t. the Lebesgue measure, then:

$\mu_\omega$  is a random SRB measure.

- This theorem applies to a large class of dissipative stochastic systems, namely the hypoelliptic ones that exhibit a Lyapunov exponent  $> 0$ : they all support a random SRB measure.
- Furthermore, we have the important relation:

$$\mathbb{E}(\mu_\bullet) = \rho, \quad (4)$$

where  $\rho$  is the stationary solution of the Fokker-Planck equation, when the latter is unique.

## The Ruelle response formula

- Physically, the challenge is to find the trade-off between the physics present in the model and the stochastic parameterizations of the missing physics.  
From a mathematical point of view, climate sensitivity can be related to **sensitivity of SRB measures**.
- The **thermodynamic formalism à la Ruelle, in the RDS context**, helps to understand the response of **systems out-of-equilibrium**, to changes in the parameterizations (Gundlach, Kifer, Liu).
- **The Ruelle response formula**: Given an SRB measure  $\mu$  of an autonomous chaotic system  $\dot{x} = f(x)$ , an **observable**  $G : X \rightarrow \mathbb{R}$ , and a smooth time-dependent perturbation  $X_t$ , the time-dependent variations  $\delta_t \mu$  of  $\mu$  are given by:

$$\delta_t \mu(G) = \int_{-\infty}^t d\tau \int \mu(dx) X_\tau(x) \cdot \nabla_x (G \circ \varphi_{t-\tau}(x)),$$

where  $\varphi_t$  is the flow of the unperturbed system  $\dot{x} = f(x)$ .

## The susceptibility function

- In the case  $X_t(x) = \phi(t)X(x)$ , the Ruelle response formula can be written:

$$\delta_t \mu(G) = \int dt' \kappa(t - t') \phi(t'),$$

where  $\kappa$  is called the **response function**. The **Fourier transform**  $\hat{\kappa}$  of the response function is called the **susceptibility function**.

- In this case  $\delta_t \mu(\hat{G})(\xi) = \hat{\kappa}(\xi) \hat{\phi}(\xi)$  and since the r.h.s. is a product, there are no frequencies in the linear response that are not present in the signal.
- In general, the situation can be more complicated and the theory gives the following criterion of high sensitivity:  
**ℳ: Poles of the susceptibility function  $\hat{\kappa}(\xi)$  in the upper-half plane**  
**⇒ High sensitivity of the system's response function  $\kappa(t)$ .**
- RDS theory offers a path for extending this criterion when random perturbations are considered.



# Climatic uncertainties & moral dilemmas



**Thought leaders**  
Rice, top left, spoke of multilateralism, while Bono, left, demanded more action on poverty. Presidents Karzai and Musharraf, right, both face troubles at home

♥ ... keep today's climate for tomorrow?



**Agitator Gore**  
wants a global compact to tackle climate change and poverty

♥ **Feed the world today or...**

Davos, Feb. 2008, photos by *TIME Magazine*, 11 Feb. '08;  
see also Hillerbrand & Ghil, *Physica D*, 2008, **237**, 2132–2138,  
[doi:10.1016/j.physd.2008.02.015](https://doi.org/10.1016/j.physd.2008.02.015) .



# The Biofuel Myth

- ◆ Fine illustration of the moral dilemmas (\*).
- ◆ Conclusion: “**festina lentae**,” as the Romans (\*\*) used to say..

(\*\*) ~ Han dynasty

(\*) Hillerbrand & Ghil, *Physica D*, 2008, [doi:10.1016/j.physd.2008.02.015](https://doi.org/10.1016/j.physd.2008.02.015), available on line.





# Climate Change 1816–2008



T. Géricault, 1819,  
Le Louvre

M. Gillot, 2008,  
Le Monde

